


***TRANSURETHRAL  
RESECTION OF A  
BLADDER TUMOUR  
(TURBT)  
PATIENT  
INFORMATION***

The information contained in this booklet is intended to assist you in understanding your proposed surgery. Some of the information may or may not apply to you. Please bring this book with you to hospital as it is a useful guide. Feel free to discuss any issues and questions you may have about your surgery with the medical and nursing staff looking after you.

## ***What does the Bladder do?***

The bladder is a hollow, muscular organ in the pelvis behind the pubic bone.

The function of the bladder is to collect, store and expel urine as the kidneys produce it. When the bladder is full, the nerves that supply it send a message to the brain that you need to pass urine. Then, under your control, the outlet pipe (urethra) muscles relax and the bladder contracts until it is empty of urine.



# ***What is a Transurethral Resection of a Bladder Tumour?***

A Transurethral Resection of a Bladder Tumour (TURBT) is the removal of abnormal tissue (tumour) from the bladder. The tissue is removed by passing a special instrument up the urethra. This instrument has a light, a telescope for viewing the bladder, and a special electrode that cuts away the bladder tissue.

Occasionally bladder tumours are benign (non-cancerous) but usually they are malignant (cancerous). There are several different types of bladder cancer: the most common is transitional cell carcinoma (TCC). Other less common tumours are squamous cell carcinoma (SCC), adenocarcinoma and sarcoma.

Factors that may increase the likelihood of developing bladder cancer include smoking, exposure to chemicals, and infection from some tropical parasites.

The treatment of bladder cancer depends on the type of tumour and how far it has spread. Your doctor will discuss the type of tumour you have and the likelihood of it spreading. The doctor will also discuss the ongoing management of it with you.

## ***Potential Complications***

This procedure carries a small risk of excessive bleeding and urinary tract infection. You will be monitored for these risks and treated promptly if they occur.

- **Excessive bleeding**

Your vital signs (blood pressure and pulse) and urine will be monitored for signs of excessive bleeding.

- **Infection**

Your temperature will be monitored for early signs of infection and intervention will be put in place if it occurs. To reduce the risk of infection antibiotics are given directly into your bloodstream during your operation and continued post-operatively if necessary. You can also assist with the prevention of infection by maintaining good hygiene and doing your deep breathing exercises. Early mobilisation also helps.

- **Bladder perforation (hole)**

Due to the nature of this operation, it is sometimes necessary to cut deeply in order to remove the bladder tumour. This may produce a small hole in the muscle of the bladder wall. If this happens, a drainage tube (catheter) may be left in the bladder for a few extra days to rest the bladder so that the area is able to heal.

## ***Length of Stay***

The usual length of stay is one to two days. However, if other procedures are required it may be necessary for you to remain in hospital for a few more days. Your doctor will discuss this with you.

## ***Before Surgery***

### **Informed consent**

After consultation with the doctor you will be asked to sign a form to give written consent for the surgeon to perform the operation and for an anaesthetic to be administered. Relevant sections of the form must also be completed if you agree to a blood transfusion and/or if your particular surgery involves the removal of a body part and you wish to have this returned. Our

expectation is that you feel fully informed about all aspects of your surgery before giving written consent. The following health professionals are available to help you with this process.

## **Nurses**

A nurse will explain what to expect before and after surgery. Please ask questions and express your concerns; your family or people close to you are welcome to be involved.

When you are discharged from hospital your nurse will arrange for you to receive ongoing support, advice and practical help, if needed.

## **Tests**

### **Blood samples**

Samples of your blood will go to the laboratory to check your general health before surgery.

### **Blood transfusions**


A sample of your blood will go to the blood bank to identify your blood type so this can be matched with donated blood. This donated blood is then ready for transfusion during or after surgery if required. **We will need your written consent before a transfusion is able to take place.**

### **Midstream urine**

A sample of your urine is sent to the laboratory to check that there is no bacteria.

### **Chest x-ray**

If requested by the doctor or anaesthetist, a Chest x-ray will be performed to check on the health of your lungs.



## **ECG**

An electrocardiogram (ECG) of your heart may be required depending on your age and any diagnosed heart conditions.

## **Other measures**

### **Nil by mouth**

As your stomach should be empty before an anaesthetic, you must not eat anything or drink milk products six hours prior to surgery. You may, however, be able to drink clear fluids up to two hours before surgery - the Pre-Admission Clinic or ward nurse will clarify this with you.



## ***Wound site - What to expect***

Although there is no visible wound, there will be a wound inside your bladder which you may be aware of when you begin to pass urine again. The raw area where tissue has been removed from your bladder lining may cause some discomfort ie. burning or stinging when your bladder fills and/or you pass urine. You will be offered some medications to relieve any discomfort experienced.

## ***After Surgery***

You are transferred to the Recovery Room next to the theatre. Your condition is monitored and when you are awake and comfortable a nurse and an orderly will escort you back to the ward on your bed.

## **On the ward**

**Your nurse will check the following regularly:**

- Vital signs - your blood pressure, pulse, respiration rate and temperature
- The severity and location of any pain or discomfort
- The effectiveness of pain relief
- The colour of urine you are producing
- The amount of oxygen in your blood

## **You may have Intravenous fluids**


A small tube (leur) is placed into a vein in the forearm to give you fluids and medications.

## **Oxygen**

Oxygen is often given for the first 24 hours after surgery via nasal prongs or a facemask to help with breathing and healing.

## **Continous bladder irrigation (Through & through)**

Your urine will contain some blood after surgery that can clot and cause blockages. Therefore, for 6-24 hours your catheter is connected to an irrigation system that flushes the bladder in order to prevent these problems. If a blockage does occur, your nurse will flush the catheter with a syringe filled with salt water (saline) in order to unblock it.



## **Urinary catheter (IDC)**

You will have a tube in your bladder that will drain the urine (urethral catheter). This can be secured to your leg for comfort.

## **Pain relief after your surgery**

Your nurse will work alongside your doctors and the anaesthetist to keep your pain at a minimum.

The **PAIN SCORE** is a way of your nurse establishing how much pain you are experiencing by asking you to grade your pain from 0 to 10 where 0 = no pain and 10 = the worst pain you can imagine.

The following methods of pain relief may be used singly or in combination with each other.

### **Intravenous (IV) pain relief**

Intravenous pain relief can be administered if required.

### **Oral pain relief**

When you are able to drink, you may have tablets by mouth (orally).



### **Urinary alkalisation**

As mentioned, you may feel a burning sensation when you pass urine. Your nurse can provide you with sachets of urinary alkalisers (eg. Citravescent or Ural) to help with this. This is a pleasant tasting powder that is added to the fluid you drink. It makes the urine less “acid” and therefore less likely to sting.

## **Food and fluids**

You will be able to progress from sips to a full diet in a short space of time after you have fully woken up from your anaesthetic. Medications are available for the relief of nausea and vomiting, if they occur.

## **Mobility**

Your movement will be restricted for as long as the continuous bladder irrigation remains attached to your catheter. You will be able to be up and about when this has been removed. Your mobility will increase as you recover; mobilisation will assist your recovery.



## ***Removal of Drips and Drains***

### **Intravenous fluid (IV)**

The IV fluid is usually stopped the day of surgery or the day after providing the amount of blood in the urine is acceptable. The leuc (plastic tube) is removed when you no longer require intravenous medications.

### **Urinary catheter**

The urinary catheter is usually removed one to two days after surgery. Sometimes after the catheter has been removed there is temporary difficulty with control of your urine flow. This should settle down as the bladder heals. Occasionally people are unable to pass urine naturally because of swelling or the presence of blood clots in the bladder. If this occurs, a nurse will reinsert a

urinary catheter into your bladder. This will allow the swelling to reduce or enable the flushing and removal of any blood clots.

## **Important information for when your catheter is removed**

- It is important to try to drink at least two to three litres of fluid a day to aid the flushing of any blood that remains in your urine. This is easier if you vary your fluids eg. fruit juice, cordial, tea, etc., in addition to water. Do not drink more than this.
- Drink small amounts regularly eg. one to two glasses over each hour. Drinking large amounts at once may make you feel bloated or nauseated.
- Go to the toilet when you get the desire – don't strain to pass urine.
- Use a new bottle each time you pass urine – this allows your nurse to check that you are emptying your bladder properly.
- Initially it may burn when you pass urine and you may pass urine frequently. This usually improves over the following days and can be relieved by drinking fluids as discussed previously.
- Please inform your nurse if any of the following occur:
  - you are unable to pass urine despite having the urge to go
  - you have pain or discomfort in your lower abdomen (stomach)
  - you have pain at the tip of your penisThese symptoms could indicate difficulty with emptying your bladder. Your nurse will be able to assist you.
- The nurse will use a bladder scanner (small, painless ultrasound) to check if you are emptying your bladder properly.
- If you have not moved your bowels since your operation, please tell your nurse.
- Approximately 10-14 days after surgery you may pass slightly bloodstained urine again. This is normal and should stop within a day or two – continue to drink plenty.

**Contact your GP if blood clots appear, if bleeding is heavy or you have any problems.**

## ***Discharge Advice***

- Your hospital doctor will provide your first sickness benefit certificate/medical certificate and will advise you when to return to work.
- There remains a risk of bleeding for several weeks after TURBT surgery. This means that your urine may have a pinkish tinge for up to three weeks. This will settle as your body heals. Meanwhile, it is important to continue to drink two to three litres of fluid a day to maintain flushing of your bladder.
- Avoid heavy lifting or strenuous activity for at least four to six weeks - contact sports are not generally recommended. Sexual activity may be resumed at this time or when you feel comfortable to do so.
- Maintain a regular bowel habit and avoid constipation as straining to pass a bowel motion may cause more blood in the urine.
- If bleeding occurs and your urine flow is obstructed altogether it is important to return to hospital quickly so that it can be attended to.
- If you experience chills, fever or pain in your bladder or back, or if your urine is cloudy and smells offensive, then see your GP promptly.

# ***Follow-up***

## **Discharge letter**

You and your GP will receive a copy of a letter outlining the treatment you received during your stay in hospital. This will be sent to you if it is not completed by the time you leave hospital.

## **General Practitioner (Family doctor)**

When you are discharged from hospital you will be under the care of your family doctor who will look after your general health and monitor any problems you may have.

## **Outpatients appointments**

After a Transurethral Resection of a Bladder Tumour (TURBT) you will have your bladder checked regularly in the Outpatients Department. Results from the surgery will be discussed with you at this time or at a separate clinic appointment. Details of this appointment will be sent to you.



3 References: Mosby's Genitourinary Disorders, Clinical Nursing, Mikel Gray 1992  
Urological Nursing 3rd Edition, Urological Nursing' 2004  
Campbell's Urology 7th Edition, Urology, 1998