

Vaginal bleeding and/or pain in early pregnancy in the haemodynamically stable patient

Overview

Purpose

To ensure that women presenting with vaginal bleeding and/or pain in early pregnancy receive the best available evidence-based care and treatment.

Scope

Nursing, Medical and Clerical staff caring for women presenting with vaginal bleeding and/or pain in early pregnancy

References


1	Australian Society for Ultrasound in Medicine (2012). <i>Guidelines for the Performance of First Trimester Ultrasound</i> . New South Wales. Australian Society for Ultrasound in Medicine.
2	Royal College of Obstetricians and Gynaecologists (RCOG) (2006). <i>The Management of Early Pregnancy Loss. Guideline No. 25</i> . United Kingdom. Royal College of Obstetricians and Gynaecologists.
3	Royal College of Obstetricians and Gynaecologists (RCOG) (2011). <i>The Management of Early Pregnancy Loss. Guideline No. 25. Appendix October 2011</i> . United Kingdom. Royal College of Obstetricians and Gynaecologists.
4	Waitemata District Health Board. (2013). <i>Pregnancy of Unknown Location</i> . Auckland: Waitemata District Health Board.

Issued by	Clinical Nurse Specialist Early Pregnancy Clinic	Issued Date	June 2021	Classification	010210-22-003
Authorised by	Clinical Director Gynaecology	Review Period	36 months	Page	1 of 2

This information is correct at date of issue. Always check on Waitemata DHB Controlled Documents site that this is the most recent version.

Vaginal bleeding and/or pain in early pregnancy in the haemodynamically stable patient

Flowchart



Issued by	Clinical Nurse Specialist Early Pregnancy Clinic	Issued Date	June 2021	Classification	010210-22-003
Authorised by	Clinical Director Gynaecology	Review Period	36 months	Page	2 of 2

This information is correct at date of issue. Always check on Waitemata DHB Controlled Documents site that this is the most recent version.